

BIOSPORINE® 3

Topical Cream

Triple Topical Broad Spectrum Antibiotic Association



mms

Formulation

Each 30 g contains: Neomycin Sulfate 1.3161 mg, Polymyxin B Sulfate 0.18 mg, Zinc Bacitracin 1.62 mg Excipients c.s.p 30 g

Indications

Treatment of wounds, bacterial infections (Gram + and Gram-) of the skin and ears -caused by sensitive germs- in domestic animals. Indicated for: exudative, eczematous, seborrheic or contact dermatitis, intertrigo or other pyoderma, lacerations, abrasions, surgical wounds, adjuvant treatment in cases of allergic dermatitis due to fleas, dermatosis and alopecia, otitis externa (acute or chronic), infections of the paranal gland in canines.

Dosage and Administration

Topical route on the skin, mucous membranes, wounds or within the ear canal or paranal glands. In dermal conditions: clean and dry the application area. Apply a small amount and distribute evenly. Otic conditions: clean, use the plastic cannula and apply in the ear canal until the affected area is completed and massaged. Infected paranal gland or cystic areas: drain, clean and fill using the cannula if necessary. The frequency and duration of treatment will depend on the severity of the injury and the criteria of the Veterinarian. In most cases 1 - 4 times / day. The duration can vary between 2 to 6 weeks.

Commercial Presentation

Tube x 15g, Tube x 30g,

SENASA Registry

SENASA Perú: F.82.39.I.0421

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.