

## COMBIÓTICO L.A. SPI

### Soluble Powder

Synergistic Antibiotic Combination of Wide Spectrum and Long Action



### Formulation

Combiótico® L.A. SPI 3.6 g Bottle 1: Penicillin G Benzathine 2 000 000 I.U., dihydrostreptomycin sulfate equivalent to 2 g base. Bottle 2: Sterile diluent 8 mL. Each reconstituted mL contains: Benzathine Penicillin G 200,000 IU, dihydrostreptomycin sulfate equivalent to 0.2 g base, excipients q.s.p.1 mL. Combiótico® L.A. SPI 13.5 g Bottle 1: Penicillin G Benzathine 7,500,000 I.U., dihydrostreptomycin sulfate equivalent to 7.5 g base. Bottle 2: Sterile diluent 30 mL. Each reconstituted mL contains: Benzathine Penicillin G 200,000 IU, dihydrostreptomycin sulfate equivalent to 0.2 g base, excipients q.s.p.1 mL.

### Indications

Treatment and prevention of local or systemic infections caused by germs susceptible to penicillin and / or dihydrostreptomycin, provided that a long-acting effect is required (spaced doses)

### Dosage and Administration

Intramuscular injectable route (in intramuscular and subcutaneous canines), repeat after 72 hours if necessary. Cattle, sheep, goats, horses, South American camelids and pigs: the recommended dose is 1 mL for every 10-20 kg of live weight. In young animals: Calves, lambs, foals, piglets, the dose is 1 - 2 mL / 25 kg. Canines and felines: the recommended dose is 1 - 2 mL / 25 kg of live weight.

### Commercial Presentation

Bottle of 3.6g, Bottle of 8mL of diluent, 30mL bottle of diluent, 13.5mL bottle of diluent,

### SENASA Registry



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.