

## DOXIMUTIN® WS

### Soluble Powder

Powerful Additive-Synergistic  
Pleuromutilin-Tetracycline Antibiotic Combination



### Formulation

Each 100 g contains: Tiamulin (as fumarate) 10 g, Doxycycline (as a hyclate) 15 g Excipients c.s.p. 100g

### Indications

Treatment and prevention of respiratory, enteric, mixed and / or complicated diseases caused by those microorganisms that are sensitive to tiamulin and/or doxycycline, as well as secondary bacterial infections. Birds: chronic respiratory disease (CRD), infectious coryza, aerial sacculitis, infectious sinusitis and pneumonia. Necrotic enteritis, bacterial enteritis, colisepticemia, avian cholera, omphalitis, infectious synovitis. E. coli infections associated with rhinotracheitis in turkeys. Stress or early death from non-specific causes. Pigs: mycoplasmosis, mycoplasmal (enzootic) pneumonia, pneumonia, atrophic rhinitis, swine pleuropneumonia, Glässer disease, necrotic enteritis, porcine proliferative enteropathy (PPE), porcine ileitis, porcine dysentery, colitis, streptococcal lymphadenitis and cervical synovitis, infectious synovitis and joint abscesses. During periods of stress to maintain body weight gain.

### Dosage and Administration

Oral route mixed in the drinking water (or via food). Poultry: Preventive: 1 kg / 1,500 L of water (or 750 kg of food). Therapeutic: 1 kg / 1,000 - 1,500 L (or in 500-750 kg of food) for 3-5 days. Double the dose in cases of chronic respiratory disease. Pigs: Preventive: 1 Kg / 3,000 L of water (or in 1,500 kg of food). Therapeutic: 1 Kg / 2,000 - 3,000 L of water (or 1,000-1,500 kg of food) for 3-7 days

### Commercial Presentation

Sachet x 100 g, Sachet x 1 kg, Sachet x 10 kg, Sachet x 15 g, Sachet x 20g,



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.

### SENASA Registry