

FLORAVIVA® 12 CEPAS SOFT CHEWS MEDIUM & LARGE

Tablet Oral

Protective symbiotic, improver and recuperator of intestinal balance Live Active Cultures - Gut Health



Formulation

Each 4 g chewable tablet contains: Probiotics (12 cultures) 20 x 108 CFU, Active Organic Dry Yeast Extract 200 mg, B-glucans 80 mg, Fructooligosaccharides (FOS) 80 mg, Mannan ligosaccharides (MOS) 80 mg, Lactoferrin 40 mg, Yucca schidigera extract 40 mg, Glutamic acid 32 mg, L-Threonine 24 mg, Vitamin A (Retinol palmitate) 10,000 IU, Vitamin D3 (as cholecalciferol) 2,000 IU, Vitamin E (as alpha tocopheryl acetate) 16 mg, Vitamin K3 (as menadione sodium bisulfite) 1,000 ug, Vitamin B1 (Thiamine hydrochloride) 16 mg, Vitamin B2 (Riboflavin 5 sodium phosphate) 16 mg, Vitamin B3 (Nicotinamide) 40 mg, Vitamin B5 (Calcium pantothenate) 60 mg, Vitamin B6 HCl (Pyridoxine) 8 mg, Vitamin B7 (Biotin) 80 ug, Vitamin B9 (folic acid) 1,000 ug, Vitamin B12 (Cyanocobalamin) 40 ug, Vitamin C (as sodium ascorbate) 100 mg, Copper (Copper Chelate-AHM2) 2,040 ug, Zinc (Zinc Chelate-AHM2) 2,584 ug, Manganese (Manganese Chelate-AHM2) 2,100 ug, L-Selenomethionine 48 ug, Methionine (AHM3) 216 mg, Excipients and flavorings q.s.ad. 4 g.

Indications

Auxiliary in the recomposition and stabilization of the intestinal flora and physiological digestion, as well as improving immunity and protecting the intestinal structure and functionality in cases of: - Diarrhea or digestive disorders, poor fecal quality (soft stools). - Acute enteritis.- Allergies or intolerance to food. - Poor fecal quality, flatulence. - Antibiotic, chemotherapy or antiparasitic therapy. - Birth and/or weaning. - Stress conditions: postoperative, childbirth, excessive heat or cold, travel, vaccination, heat, exhibitions, mating. - Infectious diseases (parasitic, viral or bacterial). Strengthens the immune system. - Changes in diet or alterations in the type of food. - Poisoning and/or poisoning. - Neonates and young growing animals. - Old age animals. - Intestinal dysbiosis (imbalance of the intestinal flora). - Pancreatic problems, malabsorption syndrome and chronic liver disorders.

Dosage and Administration

Este contenido es solo para referencia informativa y no debe ser considerado como recomendaciones de uso.

It is administered orally, according to the size of the pet. Administer 1 to 2 times a day