

HEPAVIAR® OS

Oral Solution

Hepatoprotective - Lipotropic - Detoxifying
Balanced Association



Formulation

Each 100 mL contains: L-Carnitine (as L-Carnitine hydrochloride) 5000 mg, Silimarine 2000 mg, Artichoke extract 2000 mg, DL-Methionine 2000 mg, Choline chloride 7500 mg, Betaine (as Betaine hydrochloride) 1000 mg, L-Lysine (as L-Lysine hydrochloride) 2000 mg, N-Acetyl-L-Methionine 1000 mg, Magnesium (as magnesium sulphate) 1010 mg, Sorbitol 20 000 mg, excipients q.s. ad 100 mL.

Indications

Diets with high energetic content, intoxications, treatment or prevention of fatty liver syndrome (hepatic lipidosis), treatment of mycotoxicosis, preparation of birds for forced molt, appetite and alimentary conversion improver. Acute and chronic hepatitis, colic syndrome, infectious conditions and gastrointestinal disorders. In all cases where medications that affect the hepatic parenchyma and its performance is administered (antibiotics, antiparasitary, anesthetics, tranquilizers, etc.) and in the post-operative period.

Dosage and Administration

Poultry: Preventive: In commercial layers: between weeks 28 and 40, every 4 weeks during 3 - 5 consecutive days, and in broiler chickens: 3 - 5 days with the feed change of the last day of the program: 1 L/2000 L of water. Treatment: 1 L/1000 - 1500 L of drinking water, during 5 - 7 days. Pigs: 1 L/1000 - 2000 L of water. Other especies: cattle and horses: 1 mL/5 - 10 kg of body weight during 5 - 7 days; sheep, goats, camelids, rabbits, dogs and cats: 1 mL/2.5 - 5 kg of body weight during 5 - 7 days.

Commercial Presentation

Bottle x 100 mL, Bottle x 250 mL, Bottle x 500 mL, Bottle x 1 L, Bottle x 5L, Bottle x 20L,



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.