

LIQUAMOX® C

Powder for Oral Suspension

Broad-spectrum beta-lactam antibiotic with bacterial resistance inhibitor



Formulation

Amoxicillin (as amoxicillin trihydrate) 40 mg, Clavulanic acid (as potassium clavulanate) 5.7 mg, excipients q.s.ad 1 mL

Indications

For the treatment of infections caused by susceptible strains (aerobic and anaerobic bacteria): periodontal, skin and soft tissue infections caused by susceptible species of Staphylococcus spp. E. coli, Pasteurella spp. and Streptococcus spp. Suitable for respiratory and urinary tract infections, including those caused by susceptible E. coli species. Treatment of osteomyelitis and other bacterial complications sensitive to amoxicillin.

Dosage and Administration

Preparation: shake the bottle to loosen the powder, add water to the bottle mark and shake well. After reconstitution, each mL contains 40 mg of amoxicillin (as trihydrated salt) and 5.7 mg of clavulanic acid (as potassium salt). Let stand for 5 minutes to ensure complete dispersion. Canines: 1 mL/2-3.5 kg every 8-12 hours for 7-28 days. Cats: 1 mL/2-3.5 kg every 8-12 hours for 5-14 days. The reconstituted suspension must be stored in refrigeration (between 2° C and 8° C) and can be used for up to 7 days.

Commercial Presentation

Bottle x 11 g,

SENASA Registry

SENASA Peru: F.82.32.I.0151



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.