

## NEO-TERRACICLINA® WS

### Soluble Powder

Broad-spectrum and Double Effect Antibiotic Combination



### Formulation

Each 100 g contain: Oxytetracycline 20 g, Neomycin 20 g, excipients q.s. ad 100 g.

### Indications

Treatment and prevention of infections caused by Gram-positive or Gram-negative germs sensitive to oxytetracycline and/or neomycin, mainly respiratory and gastrointestinal infections, prevention of secondary bacterial infections and prevention of early mortality. Others include diarrhea, non-specific enteritis, enterotoxemia, salmonellosis, colibacillosis, dysentery, cholera, coryza, non-specific pneumonia, atrophic rhinitis, chronic respiratory disease, airsacculitis, synovitis, omphalitis, conjunctivitis, MMA, hexamithiasis and infections secondary to viral disorders. Growth promoter.

### Dosage and Administration

By oral route mixed in drinking water (or with feed). Poultry: Promoter: 200 - 250 g/1000 kg of feed. Preventive: 250 - 500 g/1000 liters of water or 500 kg of feed. Treatment: 500 g - 1 kg/1000 liters of water or 500 kg of feed during 3 - 5 days. Pigs: Promoter: 200 - 250 g/1000 kg of feed. Preventive: 250 g/1000 liters of water or 400 kg of feed. Treatment: 500 g - 1 kg/1000 liters of water or 400 kg of feed, during 4 - 5 days. Pre-ruminants: Preventive: 60 mg of the product/kg of body weight (0.6 g/10 kg of b.w. daily); treatment: 120 mg/kg of b.w. (1.2 g/10 kg of b.w. daily) during 4 - 5 days (in water, milk or feed). Guinea pigs and rabbits: Preventive: 50 g/100 liters of water. Treatment: 100 g/100 liters of water during 3 - 5 days.

### Commercial Presentation

Jar x 1 Kg, Sachet x 15 g, Jar x 100 g,

### SENASA Registry

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.  
SENASA Perú: F.03.26.N.1806