

NEO-TERRACICLINA® WS

Soluble Powder

Broad-spectrum and Double Effect Antibiotic Combination



Formulation

Each 100 g contain: Oxytetracycline 20 g, Neomycin 20 g, excipients q.s. ad 100 g.

Indications

Treatment and prevention of infections caused by Gram-positive or Gram-negative germs sensitive to oxytetracycline and/or neomycin, mainly respiratory and gastrointestinal infections, prevention of secondary bacterial infections and prevention of early mortality. Others include diarrhea, non-specific enteritis, enterotoxemia, salmonellosis, colibacillosis, dysentery, cholera, coryza, non-specific pneumonia, atrophic rhinitis, chronic respiratory disease, airsacculitis, synovitis, omphalitis, conjunctivitis, MMA, hexamithiasis and infections secondary to viral disorders. Growth promoter.

Dosage and Administration

By oral route mixed in drinking water (or with feed). Poultry: Promoter: 200 - 250 g/1000 kg of feed. Preventive: 250 - 500 g/1000 liters of water or 500 kg of feed. Treatment: 500 g - 1 kg/1000 liters of water or 500 kg of feed during 3 - 5 days. Pigs: Promoter: 200 - 250 g/1000 kg of feed. Preventive: 250 g/1000 liters of water or 400 kg of feed. Treatment: 500 g - 1 kg/1000 liters of water or 400 kg of feed, during 4 - 5 days. Pre-ruminants: Preventive: 60 mg of the product/kg of body weight (0.6 g/10 kg of b.w. daily); treatment: 120 mg/kg of b.w. (1.2 g/10 kg of b.w. daily) during 4 - 5 days (in water, milk or feed). Guinea pigs and rabbits: Preventive: 50 g/100 liters of water. Treatment: 100 g/100 liters of water during 3 - 5 days.

Commercial Presentation

Jar x 1 Kg, Sachet x 15 g, Jar x 100 g,

SENASA Registry

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.
SENASA Perú: F.03.26.N.1806