

OPTICLOX®

Ophthalmic Ointment

Long acting antibiotic for the treatment of ocular infections



Formulation

Each 5 g syringe contains: Cloxacillin (as cloxacillin benzathine) 835 mg, Excipients q.s.ad. 5 g

Indications

Treatment of various ocular infections caused by *Staphylococcus aureus*, *Bacillus subtilis* and other bacteria susceptible to cloxacillin. Treatment of bovine keratoconjunctivitis in which the presence of susceptible bacteria (including *Moraxella bovis*) is suspected or expected.

Dosage and Administration

Opticlox® is administered by topical ocular route at a rate of (cm. of ointment per eye):
Cattle and horses: approximately 5 - 10 cm. Swine, sheep, goats, camelids: approximately 5 cm.
Dogs and cats: approximately 2 cm. Open the lower eyelid and instill a steady stream of ointment into the lower conjunctival sac. A single application is usually required, but treatment can be repeated after 48-72 hours if necessary. In animals with a single infected eye it is recommended, to avoid cross infection, to treat both eyes, treating the non-infected eye first to avoid transmission of infection. Each syringe should be used only once.

Commercial Presentation

Container x 48 syringes x 5 g,

SENASA Registry

Perú: F.082.055.I.01063



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.