

VETAMYCON 6X

Topical Gel

Antibiotic, Anti-inflammatory, Antimycotic and Antipruritic Polyvalent Association



Formulation

Beclometasona dipropionato 0.025%, Gentamicina 0.3%, Clotrimazole 1%, Lignocaína 2%.

Indications

For the treatment of bacterial (Gram + and Gram-) and fungal infections and infections of the skin and ears of domestic animals. Indicated in case of otitis externa, acute and chronic. Onychomycosis and dermal conditions of fungal etiology. Dermal conditions of bacterial etiology exudative, eczematous, seborrheic or contact dermatitis, intertrigo or other pyoderma. Dermal conditions of bacterial etiology. Exudative, eczematous, seborrheic or contact dermatitis, intertrigo or other pyoderma. Lacerations, abrasions or wounds. Infections of the paranal gland in canines.

Dosage and Administration

Topical Route. In moderate conditions: from once a day to once a week; in severe conditions, it can be applied 2-3 times a day. Minimum duration of treatment: According to the etiology and the response of the course of the disease, it can vary between 2 to 6 weeks. The frequency of administration depends on the severity of the injury and the discretion of the Veterinarian.

Commercial Presentation

Tube x 15g, Tube x 30g, Tube x 80g,

SENASA Registry

SENASA Perú: F.F9.39.I.0005



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.