

DIFLOVET® OS

Oral Solution

Second Generation Quinolone Antibiotic



Formulation

Each 1 mL contains: Difloxacin (as hydrochloride) 100 mg, excipients q.s. ad 1 mL.

Indications

For Chronic Respiratory Disease (CRD) caused by sensitive strains of *Escherichia coli* and *Mycoplasma gallisepticum* in chickens and turkeys. It is also indicated for the treatment of infections caused by *Pasteurella multocida* in turkeys. Among the susceptible agents are: *Escherichia coli*, *Mycoplasma gallisepticum*, *M. meleagridis*, *M. synoviae*, *H. paragallinarum*, *E. rhusiopathiae*, *Staphylococcus aureus*, *Chlamydia psittaci*, *Pasteurella multocida* and *Salmonella* spp.

Dosage and Administration

Poultry: the administration must be daily in drinking water at the dose of 10 mg/kg of body weight, during 5 days at least. To determinate the quantity of Diflovet® Oral every 1000 L of water, the following formula must be used: N° of animals on the breeding place x mean weight of each animal (kg) x 100/Total consumption of water at the breeding place the day before (liters).

Commercial Presentation

Bottle x 100 mL, Bottle x 250 mL, Bottle x 500 mL, Bottle x 1 L,

SENASA Registry

SENASA Perú: F.03.32.N.1180



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.