

VETOCAÍNA A2

Injectable Solution

Local anesthetic



Formulation

Lidocaine hydrochloride 20 mg, Adrenaline bitartrate 0.04 mg, excipients q.s. ad 1 mL.

Indications

As a local anesthetic by infiltration (epidural or paravertebral) in cesarean deliveries, tail docking, castration, testing and diagnosis of lameness in horses. For conductive anesthesia, nerve block (such as the intercostal, mandibular and brachial plexus nerves) and subarachnoid, intra-articular infiltration, to control clinical pain. As local anesthesia for diagnostic and exploratory purposes. The powerful vasoconstrictor action of adrenaline gives an antihemorrhagic effect at the time of surgery, further prolonging the anesthetic effect of lidocaine.

Dosage and Administration

Nerve or conduction block: diagnosis injections in horses and cows: 5 - 10 mL, anesthesia of cornual nerve of the cow (dehorning): 8 - 10 mL in each side. Infiltration anesthesia: In cases of repeated administration, the total volume administered should not exceed 0.5 mL/kg of b.w., cats: 1 mL; dogs: 1 - 2 mL, horses: 100 - 200 mL per site of surgery. Sacral anesthesia: For surgery on standing animals: Horses: 5 - 10 mL (slow administration), cows: 4 - 6 mL (low administration), sheep, goats, camelids: 2 - 3 mL (slow administration). Epidural anesthesia: For surgery on lying animals: Horses: 25 - 75 mL (slow administration), cows: 15 - 60 mL (slow administration), small animals: 1.5 mL/10 cm of spinal column length (slow administration).

Commercial Presentation

Bottle x 100 mL,

SENASA Registry

SENASA Perú: F.59.01.N.0007



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.