

# **VETOCAM® 50**

### Injectable Solution

Nonsteroidal anti-inflammatory. Injectable solution with anti-inflammatory, analgesic an antipyretic action; COX-2 selective.







#### **Formulation**

Meloxicam 5 mg, Excipientsq.s.ad. 1 mL

#### **Indications**

Bovines: Used for the reduction of clinical signs of acute respiratory infections in cattle, together with adequate antibiotic therapy. Indicated for the reduction of clinical signs of diarrhea in calves over one week old and young non-lactating calves in combination with oral rehydration therapy. For postoperative pain relief in calves caused by dehorning. Swine: Used to reduce the symptoms of lameness and inflammation present in non-infectious locomotor disorders. For the relief of post-operative pain associated with minor soft tissue surgery such as castration. Dogs: For relief of inflammation and pain present in musculoskeletal disorders in both acute and chronic. To reduce inflammation and postoperative pain after orthopedic surgeries and soft tissue. Cats: Used to reduce postoperative pain after ovarian hysterectomy and other minor soft tissue surgery.

## **Dosage and Administration**

· Catlle: 10.00 mL /100 kg, by subcutaneous or intravenous route. · Swine: For locomotor disorders: 2.0 mL / 25 kg intramuscularly. If required, a second dose 24 hours after the last application. For reduction of postoperative pain: 0.4 mL / 5 kg. · Dogs: by subcutaneous or intravenous route. For muscle-skeletal disorders 0.4 mL / 10kg. For reduction of postoperative pain, if you have a period greater than 24 hours: 0.4 mL / 10 kg. · Cats: by subcutaneous route: 0.3 mL/5 kg bw before surgery.

#### **Commercial Presentation**

Bottle x 10 mL, Bottle x 20 mL, Bottle x 50 mL, Bottle x 100 mL, Bottle x 250 mL, Bottle x 500 mL, Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.