

No- Fly

Insecticide-Neonicotinoid

Plaguicide for veterinary use

agrovetmarket s.a.

COMPOSITION

Imidacloprid	0.5 g
z-9 tricosene.....	0.10g
Denatonium benzoate.....	0.01 g
Excipientes.....q.s.ad.....	100g

INDICATIONS

No-Fly is a neonicotinoid fly killer that works by blocking postsynaptic receptors (still in "resistant" insects to traditional insecticides). This mechanism is highly effective against insects and very safe for pets and humans. It is suitable for application on surfaces and areas where insects are found; both indoors and outdoors, in the livestock sector. **No-Fly** is highly effective against the common housefly: *Musca domestica*; dung fly: *Scatophaga stercoraria*; fly blue: *Calliphora vomitoria*; stable fly *Stomoxys calcitrans*; fruit fly: *Drosophila melanogaster*; common mosquito *Culex pipiens*; Tiger mosquito *Aedes albopictus*; wasp: *Vespa vulgaris*.

USE AND HANDLING INSTRUCTIONS:

- **Spread:** Place baits spread evenly on trays, dishes, or directly on the floor at a rate of 20 g per each 10 m² preferably moistened to increase their effectiveness and especially in peripheral areas outside facilities, away from the animals, in order to attract the flies inside houses and animal pens. Prevent the product from being flattened or mounds. Reactive rewetting bait points regularly and replace as deterioration, dirt and / or consumption.
- **Spraying:** Dissolve 100 g in 100 mL water (1: 1) per 50 m². Mix well, rest for 15-30 minutes and apply spray in the area, or in a targeted manner on surfaces where insects alight. Indoors, close doors and windows, during and after application for at least 30 minutes. Ventilate the area before allowing entry.
- **Painted:** Dissolve 100 g in 75 mL water. Mix well and let the mixture stand for 20 minutes. Then apply the product by brush or brush on walls, window frames and doors, support columns, girders, beams and any other place where insects land.
- **Traps:**
 - a) Dissolve 100 g of product in 100 mL of water and empty it in plastic bags and then hang them on the places frequented by insects.
 - b) Dissolve 100 g of product in 75 mL of water to impregnate (or paint) panels of wood or cardboard with the solution and hang on places frequented by insects (highly effective attraction points).

Frequency: In all cases, apply the product in times of proliferation of insects to achieve total disinfestation. Shock action: apply every 15- 21 days; maintenance action: every 30 to 45 days. The intervals may vary depending on the level and degree of the pest.

WARNINGS / RECOMMENDATIONS

The manufacturer guarantees the quality of the product in its original container and sealed, as the efficacy of the product, as long as you give the prescribed off-label use. Do not mix with other chemical products. The product can stain, so it is advisable to make a test on a small surface area to be treated before performing painting. Use the mixture within 7-8 hours of preparation. Apply on nonporous surfaces.

MEASURES TO PROTECT THE ENVIRONMENT

To reduce the risk of poisoning to non-target species is advisable to place the baits protected to prevent access to people, children, pets and birds. Container disposal: Triple rinse system: rinse three times this container and pour the solution into the mixture and then apply it unusable by crushing or perforating it, drop it in the place allocated by local authorities for this purpose. Any container that has contained pesticides should be reused.

PRECAUTIONS:

Do not eat, drink or smoke while using the product. Keep the product in its original packaging. Do not transport or store with food and clothing. After using the product, change, wash contaminated clothing and bathe with soap and water. Wear protective clothing during handling and application to enter the treated area within 24 hours. Wear gloves. In case of poisoning call the doctor immediately or take the patient to the doctor and show the label. Symptoms of intoxication: damage to the central nervous system. Ingestion: Do not induce vomiting. Get medical

attention immediately by skin contact: Immediately remove all contaminated clothing. Wash the skin areas in contact with the pesticide, using plenty of soap and water. Inhalation: Remove the patient to a ventilated, dry and ventilated. Loosen clothing and breathing monitor. Eye contact: Rinse with fresh water for 15 minutes. If eye irritation persists see a doctor. Wash hands after handling. Never drink or induce vomiting in an unconscious person. Medical advice: Atropine, until symptoms appear atropinization. In case of emergency call the information center, Poison Control.

STORAGE:

Store in a cool, dry place away from sunlight. Store between 15 ° C and 30 ° C. Keep out of reach of children and pets.

Reg. SENASA Perú: F.07.44.I.0461, Reg. Panama: RQ-663-15.

No-Fly is a product of



agrovetmarket
animalhealth

Av. Canada 3792-3798, San Luis. Lima 30 - Peru
Tel.: (511) 2 300 300

E-mail: ventas@agrovetmarket.com - Web: www.agrovetmarket.com